

A Unique Continuation Result for Klein–Gordon Bisolutions on a 2-dimensional Cylinder

C.J. Fewster*

Department of Mathematics, University of York,
Heslington, York YO1 5DD, United Kingdom.

April 3, 1998

Abstract

We prove a novel unique continuation result for weak bisolutions to the massive Klein–Gordon equation on a 2-dimensional cylinder M . Namely, if such a bisolution vanishes in a neighbourhood of a ‘sufficiently large’ portion of a 2-dimensional surface lying parallel to the diagonal in $M \times M$, then it is (globally) translationally invariant. The proof makes use of methods drawn from Beurling’s theory of interpolation. An application of our result to quantum field theory on 2-dimensional cylinder spacetimes will appear elsewhere.

*E-mail: cjf3@york.ac.uk

1 Introduction

The term ‘unique continuation’ covers a variety of results concerning the support properties of solutions to partial differential equations (or inequalities); see, for example Chapters 8, 17 and 28 of [1]. In this short paper we will study an apparently new variation on this theme concerning weak bisolutions to the massive Klein–Gordon equation on a 2-dimensional cylinder M . We will show that if such a bisolution vanishes in a neighbourhood of a sufficiently large portion of a 2-dimensional surface lying parallel to the diagonal in $M \times M$ then it must be translationally invariant. This result was motivated by a study of quantum field theory on cylinder spacetimes [2, 3] in which it is used to deduce global information from knowledge of the expectation value of the commutator $[\hat{\phi}(x), \hat{\phi}(x')]$ of a quantum field $\hat{\phi}$ for nearby points x and x' .

To formulate our result precisely, define M to be the cylinder formed as the quotient of $\mathbb{R}^2$ by the translation $z \mapsto z + 2\pi$, where (t, z) are Cartesian coordinates on $\mathbb{R}^2$, and let P be the Klein–Gordon operator on M given by

$$P = \frac{\partial^2}{\partial t^2} - \frac{\partial^2}{\partial z^2} + \mu \quad (1.1)$$

in local (t, z) -coordinates, where μ is a real constant. By a weak P -bisolution, we mean a bidistribution $\Gamma \in \mathcal{D}'(M \times M)$ such that

$$\Gamma(Pf \otimes g) = \Gamma(f \otimes Pg) = 0 \quad (1.2)$$

for all test functions $f, g \in \mathcal{D}(M)$. Such bisolutions play a well known and important rôle in linear scalar quantum field theory as ‘two-point functions’ [4]. We shall prove that weak P -bisolutions exhibit a unique continuation property modulo translational invariance across 2-dimensional surfaces in $M \times M$ of form

$$S_{pp'} = \{(T_{(\tau, \zeta)}p, T_{(\tau, \zeta)}p') \in M \times M \mid \tau \in [0, \pi], \quad \zeta \in [0, 2\pi]\} \quad (1.3)$$

for some fixed $p, p' \in M$, where $T_{(\tau, \zeta)}$ translates points by

$$T_{(\tau, \zeta)}q(t, z) = q(t + \tau, z + \zeta) \quad (1.4)$$

and $q : \mathbb{R}^2 \rightarrow M$ is the defining quotient map. These compact surfaces lie within (2-dimensional) planes parallel to the diagonal in the product manifold $M \times M$. Our result is the following.

Theorem 1.1 *Let μ be any real number other than a negative integer or zero and define P by Eq. (1.1). If Γ is a weak P -bisolution vanishing on an open neighbourhood of some surface $S_{pp'}$ of the form (1.3) then Γ is translationally invariant.*

Here, Γ is said to be translationally invariant if $\Gamma(T_{(\tau, \zeta)}f \otimes T_{(\tau, \zeta)}g) = \Gamma(f \otimes g)$ for all test functions $f, g \in \mathcal{D}(M)$ and all $(\tau, \zeta) \in \mathbb{R}^2$, where the translation $T_{(\tau, \zeta)}$ acts on test functions by

$$((T_{(\tau, \zeta)}f) \circ q)(t, z) = (f \circ q)(t - \tau, z - \zeta). \quad (1.5)$$

Theorem 1.1 could also be stated by saying that the restriction of a weak P -bisolution Γ to any neighbourhood of $S_{pp'}$ determines Γ uniquely up to the addition of a translationally invariant bisolution. We remark that nontrivial translationally invariant bisolutions which vanish near $S_{pp'}$ are easily constructed from any weak solution φ vanishing near $q(0,0)$ by writing (rather loosely)

$$\Gamma(t, z; t', z') = \varphi(t - t' + \tau, z - z' + \zeta) \quad (1.6)$$

where τ, ζ are chosen so that $p' = T_{(\tau, \zeta)}p$. In the hypotheses of Theorem 1.1 the condition $\mu \neq 0$ is essential, while the requirement that μ should not equal a negative integer is primarily a technical convenience and can doubtless be removed. To see that the result fails if $\mu = 0$, it suffices to note that if $G \in \mathcal{D}'(\mathbb{R} \times \mathbb{R})$ is a doubly 2π -periodic distribution, then (informally)

$$\Gamma(t, z; t', z') = G(t - z, t' - z') \quad (1.7)$$

is a weak P -bisolution in this case. The requirement that Γ vanish in a neighbourhood of some surface $S_{pp'}$ of the form (1.3) restricts $G(u, u')$ only in the vicinity of some diagonal $u = u' + \text{const}$, so one may easily construct nontranslationally invariant bisolutions Γ vanishing near $S_{pp'}$ if $\mu = 0$.

We will prove Theorem 1.1 in Section 3 using a result on exponential series established in Section 2 which draws on Beurling's theory of interpolation [5]. (These ideas also underpin Shannon-type theorems in signal processing (see, e.g., [6].) As mentioned above, an application of Theorem 1.1 to quantum field theory on cylinder spacetimes will appear elsewhere [2, 3].

2 A Result on Exponential Series

In this section, we use ideas and methods drawn from Beurling's theory of interpolation [5] to obtain a result on exponential series which will be the key to the continuation result proved in the next section.

Let $\Lambda \subset \mathbb{C}$ be a countable set with at most finitely many non-real elements. For each complex sequence $c = \{c_\lambda\}_{\lambda \in \Lambda}$ labelled by Λ and obeying

$$\sum_{\lambda \in \Lambda} |c_\lambda| < \infty, \quad (2.1)$$

i.e., $c \in \ell^1(\Lambda)$, there is a corresponding exponential series

$$\omega_c(x) = \sum_{\lambda \in \Lambda} c_\lambda e^{i\lambda x} \quad (2.2)$$

converging uniformly on compact subsets of $\mathbb{R}$ and a distribution $\omega_c \in \mathcal{D}'(\mathbb{R})$ defined by

$$\omega_c(f) = \int_{-\infty}^{\infty} dx \omega_c(x) f(x) = \sum_{\lambda \in \Lambda} c_\lambda \hat{f}(\lambda) \quad (2.3)$$

for $f \in \mathcal{D}(\mathbb{R})$. Here, the Fourier transform of f is defined by

$$\widehat{f}(\lambda) = \int_{-\infty}^{\infty} dx e^{i\lambda x} f(x). \quad (2.4)$$

It is essentially straightforward to show that the map

$$\begin{aligned} \omega : \ell^1(\Lambda) &\longrightarrow \mathcal{D}'(\mathbb{R}) \\ c &\longmapsto \omega_c \end{aligned} \quad (2.5)$$

is an injection, i.e., if $\omega_c(f) = 0$ for all $f \in \mathcal{D}(\mathbb{R})$, then $c_\lambda = 0$ for all $\lambda \in \Lambda$. In this section we will establish a condition for ω to be injective from $\ell^1(\Lambda)$ to $\mathcal{D}'(E)$ for E a finite interval in $\mathbb{R}$. We will require the following:

Definition 2.1 *A (countable) set $\Lambda \subset \mathbb{C}$ is said to be uniformly discrete if there exists $\delta > 0$ such that $|\lambda - \lambda'| > \delta$ for all pairs λ, λ' of distinct elements of Λ . If Λ is uniformly discrete and contains at most finitely many non-real elements, then we define the upper uniform density $D^+(\Lambda)$ of Λ to be*

$$D^+(\Lambda) = \lim_{r \rightarrow \infty} \frac{N(r)}{r} \quad (2.6)$$

where $N(r)$ is the maximum number of elements of Λ contained in any interval of form $[x, x + r]$, $x \in \mathbb{R}$.

We remark that $N(r)/r$ has a limit as $r \rightarrow \infty$ because $N(r)$ is subadditive and $N(r)/r$ is bounded above (by uniform discreteness).

Our main result in this section is based on the following, which is a slight modification of a result due to Beurling in [5].

Proposition 2.2 *If $\Lambda \subset \mathbb{C}$ is uniformly discrete and has at most finitely many non-real elements and E is any real interval of length greater than $2\pi D^+(\Lambda)$, then there exist test functions $\{f_\lambda\}_{\lambda \in \Lambda}$ in $\mathcal{D}(E)$ such that*

$$\widehat{f}_\lambda(\lambda') = \begin{cases} 1 & \lambda = \lambda' \\ 0 & \lambda \neq \lambda' \end{cases} \quad (2.7)$$

for $\lambda, \lambda' \in \Lambda$.

In the case $\Lambda \subset \mathbb{R}$, this follows directly from the proof of Theorem 1 in [5]. In that argument, Beurling constructs analytic functions $\widehat{f}_\lambda$ [which he denotes g_λ] obeying Eq. (2.7) and which are of the correct exponential type to be Fourier transforms of functions supported in E . By construction, these functions are in fact Fourier transforms of functions in $\mathcal{D}(E)$, although continuity suffices for Beurling's purposes. The result is easily extended to include finitely many non-real elements using the ideas of Lemma 4 in [5]. Beurling's result has the following immediate consequence:

Corollary 2.3 *If Λ and E obey the hypotheses of Theorem 2.2 then ω is injective as a map from $\ell^1(\Lambda)$ to $\mathcal{D}'(E)$.*

Proof: Applying Proposition 2.2, we have $c_\lambda = \omega_c(f_\lambda)$ for all $\lambda \in \Lambda$. ■

Our main result here is a slight extension of Proposition 2.2 to address more general countable sets Λ which may contain accumulation points.

Theorem 2.4 *Suppose $\Lambda \subset \mathbb{C}$ is countable with at most finitely many non-real elements. If there exists $R > 0$ such that $\Lambda_R = \{\lambda \in \Lambda \mid |\lambda| > R\}$ is uniformly discrete then $\omega : \ell^1(\Lambda) \rightarrow \mathcal{D}'(E)$ is injective whenever E is an interval of length greater than $2\pi D^+(\Lambda_R)$.*

Proof: Let a and b be any real numbers such that $a > b > 2\pi D^+(\Lambda_R)$ and suppose a sequence $c = \{c_\lambda\}_{\lambda \in \Lambda}$ exists such that $\omega_c(f) = 0$ for all $f \in \mathcal{D}(0, a)$. We will show that $c_\lambda = 0$ for all $\lambda \in \Lambda$, thus establishing that ω is injective from $\ell^1(\Lambda)$ to $\mathcal{D}'(0, a)$. Let

$$\varphi(x) = \sum_{\lambda \in \Lambda_R} c_\lambda e^{i\lambda x} \quad \text{and} \quad \psi(x) = \sum_{\lambda \in \Lambda \setminus \Lambda_R} c_\lambda e^{i\lambda x} \quad (2.8)$$

so that $\omega_c = \varphi + \psi$.

For each $\mu \in \Lambda \setminus \Lambda_R$, we apply Proposition 2.2 to the set $\Lambda_R \cup \{\mu\}$ and the interval $(0, b)$ to obtain a test function $f_\mu \in \mathcal{D}(0, b)$ such that $\widehat{f_\mu}(z)$ vanishes for $z \in \Lambda_R$ and equals unity for $z = \mu$. Now consider the 1-parameter family of test functions $(T_\alpha f_\mu)(x) = f_\mu(x - \alpha)$ in $\mathcal{D}(0, a)$ for $\alpha \in [0, a - b]$. We have $\omega_c(T_\alpha f_\mu) = 0$ by hypothesis on ω_c ; moreover

$$\varphi(T_\alpha f_\mu) = \sum_{\lambda \in \Lambda_R} c_\lambda e^{i\lambda \alpha} \widehat{f_\mu}(\lambda) = 0 \quad (2.9)$$

by definition of f_μ . Thus

$$\psi(T_\alpha f_\mu) = \sum_{\lambda \in \Lambda \setminus \Lambda_R} e^{i\lambda \alpha} c_\lambda \widehat{f_\mu}(\lambda), \quad (2.10)$$

vanishes for all $\alpha \in [0, a - b]$. Since the right-hand side of Eq. (2.10) extends to an analytic function of α [because $\Lambda \setminus \Lambda_R$ is a bounded subset of $\mathbb{C}$] we deduce that it must vanish identically. It follows that $c_\lambda \widehat{f_\mu}(\lambda) = 0$ for all $\lambda \in \Lambda \setminus \Lambda_R$ and in particular that $c_\mu = 0$ [because $\widehat{f_\mu}(\mu) = 1$].

We have now shown that $c_\mu = 0$ for all $\mu \in \Lambda \setminus \Lambda_R$. To conclude the proof, we apply Corollary 2.3 to the set Λ_R and the interval $(0, a)$ to show that $c_\lambda = 0$ for all $\lambda \in \Lambda_R$. Thus ω is injective as a map from $\ell^1(\Lambda)$ to $\mathcal{D}'(0, a)$ and hence from $\ell^1(\Lambda)$ to $\mathcal{D}'(E)$ whenever E is a real interval of length greater than $2\pi D^+(\Lambda_R)$. ■

3 Proof of Theorem 1.1

We begin with some preliminaries. Let μ be any real number other than zero or a negative integer, and let $\xi_{n\epsilon}$ (for $n \in \mathbb{Z}$, $\epsilon = \pm$) denote the distribution acting on $f \in \mathcal{D}(M)$ by

$$\xi_{n\epsilon}(f) = \int_{\mathbb{R} \times [0, 2\pi]} dt dz e^{-i\epsilon\omega_n t + inz} (f \circ q)(t, z) \quad (3.11)$$

where q is the quotient map defining M and ω_n is given by

$$\omega_n^{1/2} = \begin{cases} |n^2 + \mu|^{1/4} & n^2 > -\mu \\ e^{i\pi/4} |n^2 + \mu|^{1/4} & n^2 < -\mu. \end{cases} \quad (3.12)$$

If $\mu < 0$, finitely many of the ω_n can be imaginary. Equivalently,

$$\xi_{n\epsilon}(f) = \widehat{f}(\epsilon\omega_n, n) \quad (3.13)$$

where the Fourier transform $\widehat{f}$ of $f \in \mathcal{D}(M)$ is

$$\widehat{f}(\omega, n) = \int_{\mathbb{R} \times [0, 2\pi]} dt dz e^{-i\omega t + inz} (f \circ q)(t, z). \quad (3.14)$$

Each $\xi_{n\epsilon}$ is a weak P -solution for the corresponding value of μ ; moreover, any weak P -bisolution Γ may be expanded in terms of the $\xi_{n\epsilon}$:

$$\Gamma = \sum_{\substack{n, n' \in \mathbb{Z} \\ \epsilon, \epsilon' = \pm}} \gamma_{nn'}^{\epsilon\epsilon'} \xi_{n\epsilon} \otimes \xi_{n'\epsilon'}, \quad (3.15)$$

where the coefficients $\gamma_{nn'}^{\epsilon\epsilon'}$ grow no faster than polynomially in n, n' , so the sum converges in the weak-* topology on $\mathcal{D}'(M \times M)$. Clearly,

$$\Gamma(T_{(\tau, \zeta)} f_1 \otimes T_{(\tau, \zeta)} f_2) = \sum_{\substack{n, n' \in \mathbb{Z} \\ \epsilon, \epsilon' = \pm}} \gamma_{nn'}^{\epsilon\epsilon'} e^{i(n+n')\zeta - i(\epsilon\omega_n + \epsilon'\omega_{n'})\tau} \xi_{n\epsilon}(f_1) \xi_{n'\epsilon'}(f_2), \quad (3.16)$$

from which it follows that Γ is translationally invariant if and only if the only nonzero coefficients $\gamma_{nn'}^{\epsilon\epsilon'}$ are those on the anti-diagonal, with $n = -n'$, $\epsilon = -\epsilon'$.

We now prove Theorem 1.1, which states that a weak P -bisolution is translationally invariant if it vanishes on a neighbourhood of a diagonal set $S_{pp'}$ of the form (1.3) for $p, p' \in M$.

Proof of Theorem 1.1: We express Γ in the form (3.15) and show that all coefficients $\gamma_{nn'}^{\epsilon\epsilon'}$ other than those on the anti-diagonal must vanish if Γ vanishes in a neighbourhood N of a diagonal set $S_{pp'}$. First note that, by compactness of S , there are open neighbourhoods $\mathcal{O}_1$ of $q(t_1, z_1)$ and $\mathcal{O}_2$ of $q(t_2, z_2)$ in M such that N contains the translates $T_{(\tau, \zeta)} \mathcal{O}_1 \times$

$T_{(\tau,\zeta)}\mathcal{O}_2$ for $\tau \in E$, $\zeta \in [0, 2\pi]$ where E is a real interval of length greater than π . Thus, fixing $f_i \in \mathcal{D}(\mathcal{O}_i)$ we have

$$F(\tau, \zeta) \stackrel{\text{def}}{=} \Gamma(T_{(\tau,\zeta)}f_1 \otimes T_{(\tau,\zeta)}f_2) = 0 \quad (3.17)$$

for all τ, ζ in these ranges. If we now take Fourier components of F , defining $F_N(\tau)$ ($N \in \mathbb{Z}$) by

$$F_N(\tau) = \frac{1}{2\pi} \int_0^{2\pi} d\zeta e^{-iN\zeta} F(\tau, \zeta), \quad (3.18)$$

we obtain (comparing with Eq. (3.16))

$$\begin{aligned} F_N(\tau) &= \sum_{(n,n') \in I_N^+} \left(e^{-i\alpha(n,n')\tau} p_{nn'}^+(f_1 \otimes f_2) + e^{i\alpha(n,n')\tau} p_{nn'}^-(f_1 \otimes f_2) \right) \\ &+ \sum_{(n,n') \in I_N} e^{-i\beta(n,n')\tau} q_{nn'}(f_1 \otimes f_2). \end{aligned} \quad (3.19)$$

In this expression, the distributions $p_{nn'}^\pm, q_{nn'} \in \mathcal{D}'(M \times M)$ are defined by

$$p_{nn'}^\epsilon = \begin{cases} \gamma_{nn'}^{\epsilon\epsilon} \xi_{n\epsilon} \otimes \xi_{n'\epsilon} + \gamma_{n'n}^{\epsilon\epsilon} \xi_{n'\epsilon} \otimes \xi_{n\epsilon} & n \neq n' \\ \gamma_{nn}^{\epsilon\epsilon} \xi_{n\epsilon} \otimes \xi_{n\epsilon} & n = n' \end{cases} \quad (3.20)$$

and

$$q_{nn'} = \gamma_{nn'}^{+-} \xi_{n+} \otimes \xi_{n'-} + \gamma_{n'n}^{+-} \xi_{n'-} \otimes \xi_{n+}, \quad (3.21)$$

the frequencies $\alpha(n, n')$ and $\beta(n, n')$ are given by

$$\alpha(n, n') = \omega_n + \omega_{n'} \quad (3.22)$$

$$\beta(n, n') = \omega_n - \omega_{n'} \quad (3.23)$$

and the summations are performed over the sets $I_N = \{(n, n') \in \mathbb{Z} \times \mathbb{Z} \mid n + n' = N\}$ and $I_N^+ = \{(n, n') \in I_N \mid n \geq n'\}$.

From Eq. (3.19) we see that each $F_N(\tau)$ is an absolutely convergent exponential series with frequencies

$$\Lambda^{(N)} = \{\pm\alpha(n, n') \mid (n, n') \in I_N^+\} \cup \{\beta(n, n') \mid (n, n') \in I_N\}, \quad (3.24)$$

of which at most finitely many are non-real. Furthermore, each $F_N(\tau)$ vanishes on a real interval of length greater than π . We will apply the results of Section 2, using the $\pm\alpha(n, n')$ as the uniformly discrete part of the spectrum. The $\beta(n, n')$ in $\Lambda^{(N)}$ belong to a compact set (which depends on N) and exhibit accumulation points.

Starting with $N \neq 0$, we note that there is exactly one term in the series (3.19) for each frequency in $\Lambda^{(N)}$. There are two cases.

Case I: $\mu > 0$. In this case $\Lambda^{(N)} \subset \mathbb{R}$. Owing to the inequalities

$$\alpha(n, n') \geq \sqrt{(n + n')^2 + 4\mu} > |n + n'| \geq |\beta(n, n')| \geq 0 \quad (3.25)$$

and the fact that $\alpha(n, N - n) = 2|n| + O(1)$ as $|n| \rightarrow \infty$, the set

$$\Lambda_{|N|}^{(N)} = \{\lambda \in \Lambda^{(N)} \mid |\lambda| > |N|\} \quad (3.26)$$

is uniformly discrete with uniform upper density $D^+(\Lambda_{|N|}^{(N)}) = \frac{1}{2}$. Theorem 2.4 immediately entails that the restriction of $F_N(\tau)$ to any real interval of length greater than π uniquely determines the coefficients in the series. Accordingly, $p_{nn'}^\pm(f_1 \otimes f_2) = 0$ for $(n, n') \in I_N^+$ and $q_{nn'}(f_1 \otimes f_2) = 0$ for $(n, n') \in I_N$.

Case II: $\mu < 0$ with $-\mu \notin \mathbb{N}$. In this case $\Lambda^{(N)}$ contains finitely many non-real frequencies arising when $\min\{|n|, |n'|\} < |\mu|^{1/2}$. For $|n - n'| > |n + n'| + 2|\mu|^{1/2}$ the inequalities

$$\alpha(n, n') > \sqrt{(n + n')^2 - 2|n + n'||\mu|^{1/2}} > |\beta(n, n')| > |n + n'| \quad (3.27)$$

hold and since $\alpha(n, N - n) = 2|n| + O(1)$ as $|n| \rightarrow \infty$ the set

$$\Lambda_R^{(N)} = \{\lambda \in \Lambda^{(N)} \mid |\lambda| > R\} \quad (3.28)$$

is uniformly discrete with $D^+(\Lambda_R^{(N)}) = \frac{1}{2}$, where $R = \sqrt{N^2 - 2|N||\mu|^{1/2}}$. As in Case I, Theorem 2.4 entails that $p_{nn'}^\pm(f_1 \otimes f_2) = 0$ for $(n, n') \in I_N^+$ and $q_{nn'}(f_1 \otimes f_2) = 0$ for $(n, n') \in I_N$.

If $N = 0$, the above analysis is modified slightly because there are infinitely many zero frequency terms (since $\beta(n, n') = 0$ for all $(n, n') \in I_0$). Nonetheless, all nonzero frequency terms must vanish by identical arguments to those above, so $p_{nn'}^\pm(f_1 \otimes f_2) = 0$ for $(n, n') \in I_0^+$. The total zero frequency coefficient also vanishes, so

$$\sum_{n \in \mathbb{Z}} q_{n-n}(f_1 \otimes f_2) = 0. \quad (3.29)$$

To summarise, letting N run through $\mathbb{Z}$, we have $p_{nn'}^\pm(f_1 \otimes f_2) = 0$ for all $n \geq n'$, and $q_{nn'}(f_1 \otimes f_2) = 0$ unless $n = -n'$, in which case Eq. (3.29) holds. Furthermore, these conclusions hold for arbitrary $f_i \in \mathcal{D}(\mathcal{O}_i)$, and we may therefore deduce that the only $\gamma_{nn'}^{\epsilon\epsilon'}$ which can fail to vanish are those with $n = -n'$, $\epsilon = -\epsilon'$ (which contribute to the zero frequency part of $F_0(\tau)$). As noted at the beginning of this section, this implies that Γ is translationally invariant. The remaining condition Eq. (3.29) simply expresses the fact that Γ must vanish near the diagonal set S . ■

4 Conclusion

In this paper we have discussed an apparently new unique continuation phenomenon for weak Klein–Gordon bisolutions on a cylinder: a bisolution vanishing in a neighbourhood of a sufficiently large diagonal set must be globally translationally invariant. It

would be interesting to see how far Theorem 1.1 can be generalised. There seem to be no essential difficulties in extending to the Klein–Gordon equation for more general constant Lorentzian metrics on M , subject to an appropriate modification of the diagonal sets $S_{pp'}$. For nonconstant metrics there are no nontrivial translationally invariant bisolutions to the corresponding Klein–Gordon equation, and we speculate that a strict unique continuation result might hold in this case. It may be that an approach based on Carleman inequalities (see e.g., Chapter 17 in [1]) would be fruitful here. Finally, one may prove analogues of our result in 2-dimensional Minkowski space [3] using more standard Fourier methods. Although a straightforward generalisation to higher dimensions is not possible [3], it would be interesting to determine whether there are nonetheless analogues of this result.

Acknowledgments: The need for Theorem 1.1 arose in joint work with Atsushi Higuchi and Bernard Kay [2, 3] and I am grateful to them for discussions on this subject. I am also grateful to Maurice Dodson for introducing me to the work of Beurling employed here. This work was supported by EPSRC Grant No. GR/K 29937 to the University of York.

References

- [1] L. Hörmander, *The analysis of linear partial differential operators, Vols I–IV*, (Springer, Berlin, 1990–94)
- [2] C.J. Fewster, “Bisolutions to the Klein–Gordon equation and quantum field theory on 2-dimensional cylinder spacetimes”, [gr-qc/9804012](#)
- [3] C.J. Fewster, A. Higuchi and B.S. Kay, “How generic is F-locality? Examples and counterexamples” In preparation
- [4] R.M. Wald, *Quantum field theory in curved spacetime and black hole thermodynamics*, (University of Chicago Press, Chicago, 1994)
- [5] A. Beurling, “Mittlag-Leffler Lectures on harmonic analysis (1977–1978)” in *The collected works of Arne Beurling, Volume 2: Harmonic analysis*, L. Carleson *et al.* (eds) (Birkhäuser, Boston, 1989)
- [6] H.J. Landau, “Sampling, data transmission and the Nyquist rate” *Proc. IEEE* **55** (1967) 1701–1706